

BINARY BELL POLYNOMIALS AND DARBOUX COVARIANT LAX PAIRS

F. LAMBERT

Theoretical Physics Division, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium

S. LEBLE

*Theoretical Physics Dept, Kaliningrad State University, Russia and
Theoretical and Mathematical Physics Dept, Technical University of Gdansk, Poland*

and J. SPRINGAEL

Theoretical Physics Division, Vrije Universiteit Brussel, Pleinlaan 2, B-1050 Brussels, Belgium

(Received 14 March, 2000; revised 12 September, 2000)

1991 *Mathematics Subject Classification.* 37K10, 37K15, 37K40.

1. Introduction. Hirota representations of soliton equations have proved very useful. They produced many of the known families of multisoliton solutions, and have often led to a disclosure of the underlying Lax systems and infinite sets of conserved quantities [1,2].

A striking feature is the ease with which direct insight can be gained into the nature of the eigenvalue problem associated with soliton equations derivable from a quadratic Hirota equation (for a single Hirota function), such as the KdV equation or the Boussinesq equation. A key element is the bilinear Bäcklund transformation (BT) which can be obtained straight away from the Hirota representation of these equations, through decoupling of a related “two field condition” by means of an appropriate constraint of minimal weight. Details of this procedure have been reported elsewhere [3,4]. The main point is that bilinear BT’s are obtained systematically, without the need of tricky “exchange formulas” [1]. They arise in the form of “ \mathcal{Y} -systems”, each equation of which belongs to a linear space spanned by a basis of binary Bell polynomials (\mathcal{Y} -polynomials) [5].

An important element is the logarithmic linearizability of \mathcal{Y} -systems, which implies that each bilinear BT can be mapped onto a corresponding linear system of Lax type. However, it turns out that these linear systems involve differential operators which, even in the simplest case, do *not* constitute a Darboux covariant Lax pair. See Chapters 2 and 3 of [6]. This fact prevents us from obtaining large classes of solutions by direct application of the powerful Darboux machinery to the systems which arise by straightforward linearization of the \mathcal{Y} -systems.

Here we present a simple scheme by means of which this difficulty can be resolved for a variety of soliton equations which admit a bilinear BT that comprises a constraint of lowest possible weight (weight 2). Darboux covariant Lax pairs for the KdV, Boussinesq and KdV₅ equations are obtained in a unified manner, by exploiting the relations between the coefficients of linear differential operators connected by a classical Darboux transformation. We show, in particular, that the concept of Darboux covariance may be regarded as an elementary tool capable of

generating families of 1 + 1 dimensional soliton systems, exhibiting their close connection within a higher dimensional hierarchy (KP).

We start our discussion by recalling the main properties of the \mathcal{Y} -polynomials (derived in [5]) and by indicating how the use of the \mathcal{Y} -basis can lead systematically from the original nonlinear partial differential equations to the associated linear systems. The example of the Lax equation is instructive since this fifth order equation cannot be derived from a single quadratic Hirota equation.

2. Binary Bell polynomials. The binary polynomials that we use are defined in terms of the exponential Bell polynomials [7]

$$Y_{mx,nt}(f) \equiv e^{-f} \partial_x^m \partial_t^n e^f \quad (1)$$

as follows:

$$\mathcal{Y}_{mx,nt}(v, w) \equiv Y_{mx,nt}(f) \Big|_{f_{px,qt} = \begin{cases} v_{px,qt} & \text{if } p+q \text{ is odd,} \\ w_{px,qt} & \text{if } p+q \text{ is even,} \end{cases}} \quad (2)$$

with the understanding that $f_{px,qt} \equiv \partial_x^p \partial_t^q f$.

They inherit the easily recognizable partitional structure of the Bell polynomials

$$\begin{aligned} \mathcal{Y}_x(v) &= v_x, & \mathcal{Y}_{2x}(v, w) &= w_{2x} + v_x^2, & \mathcal{Y}_{x,t}(v, w) &= w_{xt} + v_x v_t, \\ \mathcal{Y}_{3x}(v, w) &= v_{3x} + 3v_x w_{2x} + v_x^3, & \dots & & \end{aligned} \quad (4)$$

The link between \mathcal{Y} -polynomials and the standard Hirota expressions

$$D_x^p D_t^q G' \cdot G \equiv (\partial_x - \partial_{x'})^p (\partial_t - \partial_{t'})^q G'(x, t) G(x', t') \Big|_{x'=x, t'=t} \quad (5)$$

is given by the identity

$$\mathcal{Y}_{mx,nt}(v = \ln G'/G, w = \ln G'G) \equiv (G'G)^{-1} D_x^m D_t^n G' \cdot G. \quad (6)$$

In the particular case $G' = G$ one has

$$G^{-2} D_x^m D_t^n G \cdot G \equiv \mathcal{Y}_{mx,nt}(0, Q = 2 \ln G) \equiv \begin{cases} 0 & \text{if } m+n \text{ is odd,} \\ P_{mx,nt}(Q) & \text{if } m+n \text{ is even,} \end{cases} \quad (7)$$

the P -polynomials being characterized by an equally recognizable ‘‘even part’’ partitional structure

$$\begin{aligned} P_{2x}(Q) &= Q_{2x}, & P_{x,t}(Q) &= Q_{xt}, & P_{4x}(Q) &= Q_{4x} + 3Q_{2x}^2, \\ P_{6x}(Q) &= Q_{6x} + 15Q_{2x}Q_{4x} + 15Q_{2x}^3, & \dots & & \end{aligned} \quad (8)$$

A crucial property of the \mathcal{Y} -polynomials is related to the transformation $w = v + Q, v = \ln \psi$:

$$\mathcal{Y}_{p_x, q_t}(v, w = v + Q)|_{v=\ln \psi} = \psi^{-1} \sum_{j=0}^p \sum_{\substack{k=0 \\ (j+k)=\text{even}}}^q \binom{p}{j} \binom{q}{k} P_{j_x, k_t}(Q) \psi_{(p-j)_x, (q-k)_t}. \quad (9)$$

It should also be noticed that polynomials $\mathcal{Y}_{p_x, q_t}(v, w)$, constructed with the derivatives of dimensionless variables v and w , are homogeneous expressions of weight $p + qr$ if r stands for the dimension of t (the dimension of x is chosen equal to 1).

3. \mathcal{Y} -systems associated with sech squared soliton equations. We consider three examples of sech squared soliton equations with order ranging from 3 to 5: KdV, Boussinesq, and KdV₅.

The simplest one is the KdV equation

$$\text{KdV}(u) \equiv u_t + u_{3x} + 6uu_x = 0. \quad (10)$$

Its invariance under the scale transformation

$$x \rightarrow \lambda x, \quad t \rightarrow \lambda^3 t, \quad u \rightarrow \lambda^{-2} u, \quad (11)$$

shows that u has the dimension -2. A dimensionless field Q can be introduced by setting $u = cQ_{2x}$, with c a dimensionless parameter to be determined. The resulting equation for Q can be derived from the ‘‘potential’’ equation

$$Q_{xt} + Q_{4x} + 3cQ_{2x}^2 = 0, \quad (12)$$

which can be cast into the form

$$E(Q) \equiv P_{xt}(Q) + P_{4x}(Q) \equiv G^{-2}(D_x D_t + D_x^4)G \cdot G \Big|_{G=\exp(Q/2)} = 0 \quad (13)$$

by setting $c = 1$.

The well known Hirota ‘‘two field condition’’ on G and G' , to be satisfied as a differential consequence of a bilinear BT (that one must find) takes the form [1]

$$G'^{-2}(D_x D_t + D_x^4)G' \cdot G' - G^{-2}(D_x D_t + D_x^4)G \cdot G = 0. \quad (14)$$

It corresponds to the following condition on $Q = 2 \ln G = w - v$ and $Q' = 2 \ln G' = w + v$:

$$\begin{aligned} E(w + v) - E(w - v) &= 2(v_{xt} + v_{4x} + 6v_{2x}w_{2x}) \\ &\equiv 2\{\partial_x[\mathcal{Y}_t(v) + \mathcal{Y}_{3x}(v, w)] + 6W[\mathcal{Y}_{2x}(v, w), \mathcal{Y}_x(v)]\} = 0. \end{aligned} \quad (15)$$

This condition can easily be decoupled into a pair of equations in the form of linear combinations of \mathcal{Y} -polynomials set equal to zero. It suffices to impose such a constraint on v and w ($p_j, q_j = \text{integer or zero}$, $c_j = \text{constant}$)

$$\sum_j c_j \mathcal{Y}_{p_j x, q_j t}(v, w) = 0, \quad (16)$$

of lowest possible order (or weight).

The simplest possible choice is a constraint of weight 2

$$\mathcal{Y}_{2x}(v, w) \equiv w_{2x} + v_x^2 = 0. \quad (17)$$

In order to obtain a parameter dependent decomposition we rather impose the condition

$$\mathcal{Y}_{2x}(v, w) = \lambda, \quad \lambda = \text{arbitrary parameter of weight 2}. \quad (18)$$

It leads us to the following \mathcal{Y} -system

$$\begin{aligned} \mathcal{Y}_{2x}(v, w) - \lambda &= 0, \\ \mathcal{Y}_t(v) + \mathcal{Y}_{3x}(v, w) + 3\lambda\mathcal{Y}_x(v) &= 0, \end{aligned} \quad (19)$$

the compatibility of which is subject to that of the corresponding system for ψ (setting $w = v + Q$, $v = \ln \psi$ and using relation (9))

$$\begin{aligned} (\mathcal{L}_2 - \lambda)\psi &\equiv \psi_{2x} + (Q_{2x} - \lambda)\psi = 0, \\ (\partial_t + \mathcal{L}_3)\psi &\equiv \psi_t + \psi_{3x} + 3(Q_{2x} + \lambda)\psi_x = 0, \end{aligned} \quad (20)$$

i.e. to the (λ -independent) condition

$$(Q_{xt} + Q_{4x} + 3Q_{2x})_x \equiv \partial_x E(Q) = 0. \quad (21)$$

The bilinear equivalent of the \mathcal{Y} -system (19) is obtained by means of relation (6)

$$\begin{aligned} D_x^2 G' \cdot G &= \lambda G' G, \\ (D_t + D_x^3 + 3\lambda D_x)G' \cdot G &= 0. \end{aligned} \quad (22)$$

It is the bilinear BT for KdV proposed by Hirota [1].

A similar analysis can be applied to the Boussinesq equation

$$\text{Bq}(u) \equiv u_{2t} - u_{4x} + 3(u^2)_{2x} = 0. \quad (23)$$

It shows that this equation can be derived from a potential version, obtained by setting $u = -Q_{2x}$, which can be cast into the form

$$E(Q) \equiv P_{2t}(Q) - P_{4x}(Q) \equiv G^{-2}(D_t^2 - D_x^4)G \cdot G \Big|_{G=\exp(Q/2)} = 0. \quad (24)$$

The corresponding two field condition

$$\begin{aligned} E(Q' = w + v) - E(Q = w + v) &\equiv 2(v_{2t} - v_{4x} - 6v_{2x}w_{2x}) \\ &= -2\partial_x \mathcal{Y}_{3x}(v, w) + 2v_{2t} + 6W[\mathcal{Y}_{2x}(v, w), \mathcal{Y}_x(v)] = 0 \end{aligned} \quad (25)$$

can still be decoupled into a pair of equations of the form (16) by means of a \mathcal{Y} -constraint of weight 2 (notice that in this case the dimension of $t = 2$, so that we dispose of two \mathcal{Y} -polynomials of weight 2)

$$\mathcal{Y}_t(v) + a\mathcal{Y}_{2x}(v, w) = 0, \quad a = \text{dimensionless constant to be determined.}$$

The decoupling requires $a^2 = -3$ and produces the following parameter dependent \mathcal{Y} -system ($\lambda = \text{integration constant}$):

$$\begin{aligned} \mathcal{Y}_t + a\mathcal{Y}_{2x}(v, w) &= 0, \\ a\mathcal{Y}_{x,t}(v, w) + \mathcal{Y}_{3x}(v, w) &= \lambda. \end{aligned} \quad (a^2 = -3) \quad (26)$$

The corresponding bilinear system

$$\begin{aligned} (D_t + aD_x^2)G' \cdot G &= 0 \\ (aD_x D_t + D_x^3 - \lambda)G' \cdot G &= 0 \end{aligned} \quad (27)$$

is the bilinear BT for Boussinesq obtained by Nimmo and Freeman [8]. Its compatibility is subject to that of the linear equivalent to the system (26):

$$\begin{aligned} \psi_t + a\psi_{2x} + aQ_{2x}\psi &= 0, \\ a\psi_{x,t} + \psi_{3x} + 3Q_{2x}\psi_x + (aQ_{x,t} - \lambda)\psi &= 0, \end{aligned} \quad (a^2 = -3), \quad (28)$$

i.e. to the following potential version of the Boussinesq equation:

$$\text{PBq}(Q) \equiv (Q_{2t} - Q_{4x} - 3Q_{2x}^2)_x = 0. \quad (29)$$

We now consider the KdV₅ equation

$$\text{KdV}_5(u) \equiv u_t + u_{5x} + 10uu_{3x} + 20u_x u_{2x} + 30u^2 u_x = 0. \quad (30)$$

Setting $u = cQ_{2x}$ it is found that it can be derived from the potential equation:

$$E_c(Q) \equiv Q_{x,t} + Q_{6x} + 10cQ_{2x}Q_{4x} + 5cQ_{3x}^2 + 10c^2Q_{2x}^3 = 0. \quad (31)$$

The left hand side of this equation is homogeneous of weight 6, but there is no value of c such that it be expressible as a linear combination of the weight 6 polynomials $P_{6x}(Q)$ and $P_{x,t}(Q)$.

Setting $c = 1$, we may nevertheless consider the two field condition

$$E_1(w + v) - E_1(w - v) \equiv 2\{\partial_x[\mathcal{Y}_t(v) + \mathcal{Y}_{5x}(v, w)] + R(v, w)\} = 0, \quad (32)$$

with

$$\begin{aligned} R(v, w) = -5(v_x w_{5x} - v_{2x} w_{4x} + 6v_x w_{2x} w_{3x} + 2v_x^3 w_{3x} - 3v_{2x} w_{2x}^2 \\ + 6v_x^2 v_{2x} w_{2x} + 4v_x v_{2x} v_{3x} + 2v_x^2 v_{4x} + v_x^4 v_{2x} - 2v_{2x}^3). \end{aligned} \quad (33)$$

Eliminating w_{2x} (and its derivatives) by means of the weight 2 constraint (18) it is found that the condition (32) can be decoupled into the following \mathcal{Y} -system:

$$\begin{aligned} \mathcal{Y}_{2x}(v, w) &= \lambda, \\ \mathcal{Y}(v) + \mathcal{Y}_{5x}(v, w) + 15\lambda^2 \mathcal{Y}_x(v) &= 0. \end{aligned} \quad (34)$$

Its compatibility is subject to that of the corresponding linear system:

$$\begin{aligned}\psi_{2x} + (Q_{2x} - \lambda)\psi &= 0, \\ \psi_t + \mathcal{L}_5\psi &= 0, \quad \mathcal{L}_5 = \partial_x^5 + 10Q_{2x}\partial_x^3 + 5(Q_{4x} + 3Q_{2x}^2 + 3\lambda^2)\partial_x,\end{aligned}\tag{35}$$

i.e. to the condition

$$(Q_{xt} + Q_{6x} + 10Q_{2x}Q_{4x} + 5Q_{3x}^2 + 10Q_{2x}^3)_x \equiv \partial_x E_1(Q) = 0.\tag{36}$$

4. Darboux covariant Lax pairs. Let us now go back to the KdV equation (10) and the associated linear system (20). It comprises the second order eigenvalue equation with the well known Darboux property [6], according to which (non vanishing) solutions ϕ of this equation produce gauge transformations [9]

$$G_\phi = \phi\partial_x\phi^{-1} = \partial_x - \sigma, \quad \sigma = \partial_x \ln \phi,\tag{37}$$

which map $\mathcal{L}_2(Q) = \partial_x^2 + Q_{2x}$ onto a similar operator:

$$\tilde{\mathcal{L}}_2 \equiv G_\phi\mathcal{L}_2(Q)G_\phi^{-1} \equiv \mathcal{L}_2(\tilde{Q}) \quad \text{with} \quad \tilde{Q} = Q + 2 \ln \phi.\tag{38}$$

A similar property does not hold for the evolution equation in (20). However, it is easy to see that transformations G_ϕ generated by solutions ϕ of the third order evolution equation

$$(\partial_t + L_3)\phi = 0, \quad L_3 = \partial_x^3 + b_2\partial_x + b_3,\tag{39}$$

map the operator $\partial_t + L_3$ onto the similar

$$G_\phi(\partial_t + L_3)G_\phi^{-1} = \partial_t + \tilde{L}_3, \quad \tilde{L}_3 = \partial_x^3 + \tilde{b}_2\partial_x + \tilde{b}_3,\tag{40}$$

with

$$\Delta b_2 \equiv \tilde{b}_2 - b_2 = 3\sigma_x \quad \text{and} \quad \Delta b_3 \equiv \tilde{b}_3 - b_3 = b_{2,x} + \sigma\Delta b_2 + 3\sigma_{2x}.\tag{41}$$

This follows from equation (39) and more particularly from its differential consequence

$$\sigma_t + \partial_x[\sigma_{2x} + 3\sigma\sigma_x + \sigma^3 + b_2\sigma + b_3] = 0.\tag{42}$$

The next step is to look for a system, equivalent to the system (20), in which the eigenvalue equation

$$\mathcal{L}_2(Q)\phi = \lambda\phi\tag{43}$$

is accompanied by a Darboux covariant third order evolution equation (39). The problem is to find a third order operator L_3 , with appropriate coefficients $b_2(Q)$ and $b_3(Q)$, such that $\partial_t + L_3(Q)$ be mapped, by gauge transformations (37) generated by (non vanishing) solutions ϕ of the system (43,39), onto a similar operator, $\partial_t + \tilde{L}_3(Q)$, which satisfies the covariance condition

$$\tilde{L}_3(Q) = L_3(\tilde{Q} = Q + \Delta Q), \quad \Delta Q = 2 \ln \phi. \quad (44)$$

It suffices to determine b_2 and b_3 in the form of polynomial expressions in terms of derivatives of Q (of order $r > 1$)

$$b_i = F_i(Q_{2x}, Q_{3x}, \dots), \quad i = 2, 3, \quad (45)$$

such that

$$\Delta F_i \equiv F_i(Q_{2x} + \Delta Q_{2x}, Q_{3x} + \Delta Q_{3x}, \dots) - F_i(Q_{2x}, Q_{3x}, \dots) = \Delta b_i, \quad (46)$$

with $\Delta Q_{rx} = 2(\ln \phi)_{rx}$, the Δb_i being determined by the relations (41).

Thus, in order to satisfy the first condition:

$$\Delta F_2 = F_{2, Q_{2x}} \cdot \Delta Q_{2x} + \dots = 3\sigma_x = \frac{3}{2} \Delta Q_{2x}, \quad (47)$$

one chooses

$$b_2 = F_2(Q_{2x}) \equiv \frac{3}{2} Q_{2x} + c_2, \quad c_2 = \text{arbitrary constant.} \quad (48)$$

The expression (41) for Δb_3 then becomes

$$\Delta b_3 = \frac{3}{2} Q_{3x} + 3\sigma\sigma_x + 3\sigma_{2x}, \quad (49)$$

from which Q_{3x} can be eliminated on account of equation (43)

$$Q_{3x} = -\sigma_{2x} - 2\sigma\sigma_x. \quad (50)$$

In view of the resulting expression

$$\Delta b_3 = \frac{3}{2} \sigma_{2x} = \frac{3}{4} \Delta Q_{3x}, \quad (51)$$

it is clear that the second condition

$$\Delta F_3 = F_{3, Q_{2x}} \cdot \Delta Q_{2x} + F_{3, Q_{3x}} \cdot \Delta Q_{3x} + \dots = \frac{3}{4} \Delta Q_{3x} \quad (52)$$

is satisfied if one chooses

$$b_3 = F_3(Q_{3x}) \equiv \frac{3}{4} Q_{3x} + c_3, \quad c_3 = \text{arbitrary constant.} \quad (53)$$

Setting $c_2 = c_3 = 0$ for simplicity, we find the following Darboux covariant evolution equation

$$[\partial_t + L_{3, \text{cov}}(Q)]\phi = 0, \quad L_{3, \text{cov}}(Q) = \partial_x^3 + \frac{3}{2} Q_{2x} \partial_x + \frac{3}{4} Q_{3x}. \quad (54)$$

The system (43,54) is a Darboux covariant equivalent to the former system (20) as

$$L_{3,\text{cov}}(Q) = \frac{1}{4}[\mathcal{L}_3 + 3\partial_x(\mathcal{L}_2 - \lambda)]. \quad (55)$$

The operator $L_{3,\text{cov}}(Q)$ coincides precisely to the third order operator which gives rise to the KdV equation in the Lax formalism [10]

$$[\partial_t + L_{3,\text{cov}}(Q), \mathcal{L}_2(Q)] = -\frac{1}{4}(Q_{xt} + Q_{4x} + 3Q_{2x}^2)_x. \quad (56)$$

The analogy between the Lax formalism and the requirement of Darboux covariance can be further disclosed by considering a Darboux covariant equivalent to the system (35), i.e. by looking for a system in which the eigenvalue equation (43) is accompanied by a Darboux covariant fifth order evolution equation ($c_1 = \text{constant}$)

$$[\partial_t + L_5(Q)]\phi = 0, \quad L_5(Q) = \partial_x^5 + c_1\partial_x^4 + b_2(Q)\partial_x^3 + b_3(Q)\partial_x^2 + b_4(Q)\partial_x + b_5(Q). \quad (57)$$

A transformation G_ϕ generated by a (non vanishing) solution ϕ of the system (43,57) maps $\partial_t + L_5(Q)$ onto the similar $\partial_t + \tilde{L}_5(Q)$, where

$$\tilde{L}_5(Q) = \partial_x^5 + c_1\partial_x^4 + \tilde{b}_2(Q)\partial_x^3 + \tilde{b}_3(Q)\partial_x^2 + \tilde{b}_4(Q)\partial_x + \tilde{b}_5(Q) \quad (58)$$

with

$$\Delta b_2 \equiv \tilde{b}_2 - b_2 = 5\sigma_x = \frac{5}{2}\Delta Q_{2x}, \quad (59)$$

$$\Delta b_3 \equiv \tilde{b}_3 - b_3 = b_{2,x} + \sigma\Delta b_2 + 4c_1\sigma_x + 10\sigma_{2x}, \quad (60)$$

$$\Delta b_4 \equiv \tilde{b}_4 - b_4 = b_{3,x} + \sigma\Delta b_3 + 3\sigma_x\tilde{b}_2 + 6c_1\sigma_{2x} + 10\sigma_{3x}, \quad (61)$$

$$\Delta b_5 \equiv \tilde{b}_5 - b_5 = b_{4,x} + \sigma\Delta b_4 + 2\sigma_x\tilde{b}_3 + 3\sigma_{2x}\tilde{b}_2 + 4c_1\sigma_{3x} + 5\sigma_{4x}. \quad (62)$$

These relations can again be used in a straightforward manner to determine polynomial expressions $b_i(Q)$, $i = 2, 3, 4, 5$, which satisfy the covariance requirement

$$\tilde{b}_i(Q) \equiv b_i(Q) + \Delta b_i = b_i(\tilde{Q} = Q + 2 \ln \phi). \quad (63)$$

The appropriate expression for b_2 follows immediately from relation (59) and the identity $\Delta Q_{rx} = 2(\ln \phi)_{rx}$, $r > 1$

$$b_2 = \frac{5}{2}Q_{2x} + c_2, \quad c_2 = \text{arbitrary constant}, \quad (64)$$

the subsequent $b_{i>2}$ being similarly determined (up to arbitrary constants c_i) by using relation (60–62) together with equation (50) and its differential consequences. Setting $c_4 = c_5 = 0$ and keeping c_3 as an arbitrary constant, the resulting $L_{5,\text{cov}}(Q)$ can be expressed as follows:

$$L_{5,\text{cov}}(Q) = \bar{L}_{5,\text{cov}}(Q) + c_1 \mathcal{L}_2^2(Q) + c_2 L_{3,\text{cov}}(Q) + c_3 \mathcal{L}_2(Q) \quad (65)$$

with

$$\bar{L}_{5,\text{cov}}(Q) = \partial_x^5 + \frac{5}{2} Q_{2x} \partial_x^3 + \frac{15}{4} Q_{3x} \partial_x^2 + \left(\frac{25}{8} Q_{4x} + \frac{15}{8} Q_{2x}^2 \right) \partial_x + \frac{15}{16} (Q_{5x} + 2Q_{2x} Q_{3x}). \quad (66)$$

Again it is easy to verify that the Darboux covariant system in which equation (43) is accompanied by the evolution equation

$$\phi_t + \bar{L}_{5,\text{cov}}(Q)\phi = 0 \quad (67)$$

is equivalent to the system (35) and that

$$\bar{L}_{5,\text{cov}}(Q) = \frac{1}{16} \mathcal{L}_5 + \frac{15}{16} [\partial_x^3 + (Q_{2x} + \lambda) \partial_x + Q_{3x}] [\mathcal{L}_2(Q) - \lambda], \quad (68)$$

corresponds precisely to Lax's fifth order generator of isospectral deformations of $\mathcal{L}_2(Q)$.

We notice that the appearance of the third order Darboux covariant operator $L_{3,\text{cov}}(Q)$ as part of $L_{5,\text{cov}}(Q)$ can be regarded as a direct indication of the close relationship between KdV and KdV₅ as members of the same hierarchy.

We also notice that $\mathcal{L}_2^2(Q)$ represents the actual fourth order operator $L_{4,\text{cov}}(Q)$ to be associated with $\mathcal{L}_2(Q)$ for Darboux covariance, hinting, in an elementary way, at the absence of even order flows within the KdV hierarchy.

5. Darboux covariant evolutions and KP. In contrast to the original Lax procedure, the above technique of generating soliton equations through the construction of Darboux covariant linear systems is *not* restricted to systems associated with a second order eigenvalue equation (43). It is as easy to construct a Darboux covariant system which involves a third order eigenvalue problem and a second order t -evolution (interchange of the role of \mathcal{L}_2 and L_3)

$$[\partial_t + \mathcal{L}_2(Q)]\phi = 0, \quad \mathcal{L}_2(Q) = \partial_x^2 + Q_{2x}, \quad (69)$$

$$L_3(Q)\phi \equiv \partial_x^3 + b_2(Q)\partial_x + b_3(Q) = \lambda\phi. \quad (70)$$

A gauge transformation G_ϕ generated by a (non vanishing) solution ϕ of the system (69,70) maps the operators $\partial_t + \mathcal{L}_2(Q)$ and $L_3(Q)$ onto the similar

$$G_\phi[\partial_t + \mathcal{L}_2(Q)]G_\phi^{-1} = \partial_t + \mathcal{L}_2(\tilde{Q} = Q + 2 \ln \phi), \quad (71)$$

$$G_\phi L_3(Q)G_\phi^{-1} = \tilde{L}_3(Q) \equiv \partial_x^3 + \tilde{b}_2(Q)\partial_x + \tilde{b}_3(Q), \quad (72)$$

the differences $\Delta b_i \equiv \tilde{b}_i - b_i$ being still given by relation (41). This follows again from differential consequences of equations (69,70), and, in particular from the relation

$$\sigma_t + (\sigma_x + \sigma^2)_x + Q_{3x} = 0. \quad (73)$$

On account of this relation, and the expressions (41) for Δb_2 and Δb_3 , it is clear that the covariance conditions

$$\tilde{b}_i(Q) \equiv b_i(Q) + \Delta b_i = b_i(\tilde{Q}) = Q + 2 \ln \phi), \quad i = 2, 3, \quad (74)$$

determine $b_2(Q)$ and $b_3(Q)$ up to arbitrary constants

$$b_2 = \frac{3}{2}Q_{2x} + c_2 \quad \text{and} \quad b_3 = \frac{3}{4}(Q_{3x} - Q_{xt}) + c_3. \quad (75)$$

Setting $c_2 = c_3 = 0$ it is found that $\partial_t + \mathcal{L}_2(Q)$ and

$$\widehat{L}_{3,\text{cov}}(Q) = \partial_x^3 + \frac{3}{2}Q_{2x}\partial_x + \frac{3}{4}(Q_{3x} - Q_{xt}) \quad (76)$$

constitute a Darboux covariant Lax pair for a potential version of the Boussinesq equation (23) (with a re-scaled t -variable) as

$$[\partial_t + \mathcal{L}_2(Q), \widehat{L}_{3,\text{cov}}(Q)] = -(3Q_{2t} + Q_{4x} + 3Q_{2x}^2)_x. \quad (77)$$

The closely related Darboux covariant Lax systems (43,54) and (69,70) hint at the common origin of the 1 + 1 dimensional KdV and Boussinesq equation as reductions of the 1 + 2 dimensional Kadomtsev Petviashvili equation [11]. In fact, it is natural to complete the above analysis by starting from the second order evolution equation (69) and by looking for an associated Darboux covariant t_3 -evolution

$$[\partial_{t_3} + L_3(Q)]\phi = 0, \quad L_3(Q) = \partial_x^3 + b_2(Q)\partial_x + b_3(Q). \quad (79)$$

The appropriate expressions for $b_2(Q)$ and $b_3(Q)$, are still determined by the conditions (74). In view of relation (41) and equation (73) it is clear that these expressions are given by equation (75) in which t has been replaced by t_2 .

The compatibility condition for the resulting covariant (t_2, t_3) -system (re-scaling t_3)

$$\begin{aligned} [\partial_{t_2} + \mathcal{L}_2(Q)]\phi &= 0, \\ [\partial_{t_3} + \mathcal{L}_{3,\text{cov}}(Q)]\phi &= 0, \quad \mathcal{L}_{3,\text{cov}}(Q) = 4\partial_x^3 + 6Q_{2x}\partial_x + 3(Q_{3x} - Q_{xt_2}), \end{aligned} \quad (80)$$

is subject to a condition which can be expressed as the x -derivative of an equation which involves a linear combination of P -polynomials of weight 4

$$[\partial_{t_3} + \mathcal{L}_{3,\text{cov}}(Q), \partial_{t_2} + \mathcal{L}_2(Q)] = [P_{x,t_3}(Q) + 3P_{2t_2}(Q) + P_{4x}(Q)]_x = 0, \quad (81)$$

and which does therefore correspond to a quadratic Hirota expression of degree 4

$$\mathcal{F}(D_x, D_{t_2}, D_{t_3}) = D_x D_{t_3} + 3D_{t_2}^2 + D_x^4. \quad (82)$$

Equation (81) can be regarded as a potential version of the Kadomtsev-Petviashvili equation

$$(u_{t_3} + u_{3x} + 6uu_x)_x + 3u_{2t_2} = 0, \quad (83)$$

obtained by setting $u = Q_{2x}$ and by integrating once with respect to x .

The present results suggest that a construction of higher order t_p -evolutions, $p = 4, 5, \dots$, in terms of higher operators $\mathcal{L}_{p,\text{cov}}(Q) = \partial_x^p + b_2(Q)\partial_x^{p-2} + \dots + b_p(Q)$ which are Darboux covariant with respect to $\mathcal{L}_2(Q)$, could be undertaken step by step, so as to produce higher order members of the KP hierarchy. This could be checked by identifying Jimbo and Miwa's higher degree KP equations [11] (involving higher degree Hirota forms) with potential commutator compatibility conditions between two Darboux covariant linear evolutions. In order to do so one must verify that all commutators $[\mathcal{L}_{p,\text{cov}}, \mathcal{L}_2(Q)]$ can be identified (up to differentiation with respect to x) with expressions involving linear combinations of multidimensional P -polynomials $P_{m_1 t_1, \dots, m_p t_p, \dots}(Q)$ with $t_1 = x$.

ACKNOWLEDGEMENTS. The authors wish to acknowledge the financial support extended by NATO (expert visit 975655) and that of the "Interuniversity Poles of Attraction Programme, contract nr. P4/08 — Belgian State, Prime Minister's Office - Federal Office for Scientific, Technical and Cultural Affairs".

REFERENCES.

1. R. Hirota, A new form of Bäcklund transformations and its relation to the inverse scattering problem, *Progr. Theoret. Phys.* **52** (1974), 1498–1512.
2. J. Satsuma, Higher conservation laws for the Korteweg-de Vries equation through Bäcklund transformation, *Progr. Theoret. Phys.* **52** (1974), 1396–1397.
3. F. Lambert and J. Springael, Construction of Bäcklund transformations with binary Bell polynomials, *J. Phys. Soc. Japan* **66** (1997), 2211–2213.
4. F. Lambert and J. Springael, On a direct procedure for the disclosure of Lax pairs and Bäcklund transformations, in *Chaos, Solitons and Fractals*, to appear.
5. C. Gilson, F. Lambert, J. Nimmo and R. Willox, On the combinatorics of the Hirota D-operators, *Proc. Roy. Soc. Lond. A* **431** (1996), 361–369.
6. V. B. Matveev and M. A. Salle, *Darboux transformations and solitons* (Springer-Verlag, 1991).
7. E. T. Bell, Exponential polynomials, *Ann. of Math.* **35** (1934), 258–277.
8. J. J. C. Nimmo and N. C. Freeman, A method of obtaining the N -soliton solution of the Boussinesq equation in terms of a Wronskian, *Phys. Lett. A* **95** (1983), 4–6.
9. W. Oevel and W. Schief, Darboux theorems and the KP hierarchy, in *Applications of analytic and geometric methods to nonlinear differential equations* (Kluwer Acad. Publ., Dordrecht, 1993).
10. P. D. Lax, Integrals of nonlinear equations of evolution and solitary waves, *Comm. Pure Appl. Math.* **21** (1968), 467–490.
11. M. Jimbo and T. Miwa, Solitons and infinite dimensional Lie algebras, *Publ. Res. Inst. Math. Sci. Kyoto Univ.* **19** (1983), 943–1001.